

# A systematic review and meta-analysis including GRADE qualification of the risk of surgical site infections after prophylactic negative pressure wound therapy compared with conventional dressings in clean and contaminated surgery

Fleur E.E. De Vries, MD<sup>a,\*</sup>, Elon D. Wallert, BSc<sup>a</sup>, Joseph S. Solomkin, MD<sup>b</sup>, Benedetta Allegranzi, MD<sup>c</sup>, Matthias Egger, PhD<sup>d</sup>, E. Patchen Dellinger, MD<sup>e</sup>, Marja A. Boermeester, MD<sup>a</sup>

## Abstract

**Objective:** Systematically review and Grading of Recommendations Assessment, Development, and Evaluation (GRADE) studies on prophylactic negative pressure wound therapy (pNPWT) to prevent surgical site infections (SSIs).

**Introduction:** pNPWT has been suggested as a new method to prevent wound complications, specifically SSIs, by its application on closed incisional wounds.

**Methods:** This review was conducted as part of the development of the Global Guidelines for prevention of SSIs commissioned by World Health Organization in Geneva. PubMed, Embase, CENTRAL, CINAHL, and the World Health Organization database between January 1, 1990 and October 7, 2015 were searched. Inclusion criteria were randomized controlled trials and observational studies comparing pNPWT with conventional wound dressings and reporting on the incidence of SSI. Meta-analyses were performed with a random effect model. GRADE Pro software was used to qualify the evidence.

**Results:** Nineteen articles describing 21 studies (6 randomized controlled trials and 15 observational) were included in the review. Summary estimate showed a significant benefit of pNPWT over conventional wound dressings in reducing SSIs in both randomized controlled trials and observational studies, odds ratio of 0.56 (95% confidence interval, 0.32–0.96;  $P=0.04$ ) and odds ratio of 0.30 (95% confidence interval, 0.22–0.42;  $P<0.00001$ ), respectively. This translates into lowering the SSI rate from 140 to 83 (49–135) per 1000 patients and from 106 to 34 (25–47) per 1000 patients, respectively. In stratified analyses, these results were consistent in both clean and clean-contaminated procedures and in different types of surgery, however results were no longer significant for orthopaedic/trauma surgery. The level of evidence as qualified with GRADE was however low.

**Conclusions:** Low-quality evidence indicates that prophylactic NPWT significantly reduces the risk of SSIs.

**Abbreviations:** CINAHL=Cumulative Index to Nursing and Allied Health Literature, EMBASE=Excerpta Medica Database, GRADE=Grading of Recommendations Assessment, Development, and Evaluation, MOOSE=meta-analysis of observational studies in epidemiology, NPWT=negative pressure wound therapy, OR=odds ratio, pNPWT=prophylactic negative pressure wound therapy, PRISMA=Preferred Reporting Items for Systematic Review and Meta-Analysis, RCT=randomized controlled trial, SSI=surgical site infections, WHO=World Health Organization.

**Keywords:** incisional wound therapy, prevention, prophylactic negative pressure wound therapy, surgical site infection, wound infections

Editor: Adrian Murillo.

This review was conducted in line with the development of the Global Guidelines for prevention of surgical site infections commissioned by World Health Organization in Geneva. We received no funding to perform this research.

EPD reports personal fees from Merck, Ortho-McNeil, Targanta, Schering-Plough, Astellas, Allergan, Care Fusion, Durata, Pfizer, Applied Medical, Rib-X, Affinium, Tetrphase, Televancin, R-Pharm, Cubist, Melinta, 3M and grants from Motif, outside the submitted work.

MAB reports institutional grants from J&J/Ethicon, Acclity, Ipsen, Baxter, Mylan, outside the submitted work; advisory board member of J&J/Ethicon and Acclity.

The authors have no conflicts of interest to declare.

Supplemental Digital Content is available for this article.

<sup>a</sup>Department of Surgery, Academic Medical Centre Amsterdam, The Netherlands, <sup>b</sup>Department of Surgery, University of Cincinnati College of Medicine, Cincinnati, OH, USA, <sup>c</sup>Infection Prevention and Control Global Unit,

Service Delivery and Safety, World Health Organization, Geneva, Switzerland,

<sup>d</sup>Institute of Social and Preventive medicine, University of Bern, Switzerland,

<sup>e</sup>Department of Surgery, University of Washington, Seattle, WA, USA.

\*Correspondence: Fleur E.E. De Vries, Academic Medical Centre, PO Box 22660, Amsterdam 1100 DD, The Netherlands (e-mail: f.e.devries@amc.uva.nl).

Copyright © 2016 the Author(s). Published by Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicine (2016) 95:36(e4673)

Received: 22 June 2016 / Received in final form: 27 July 2016 / Accepted: 30 July 2016

<http://dx.doi.org/10.1097/MD.0000000000004673>

## 1. Introduction

Surgical site infections (SSIs) are the number one healthcare-associated infections worldwide, with an incidence of 2% to 20%, or even higher, depending on the type of surgery and patient characteristics.<sup>[1,2]</sup> SSI are associated with increased morbidity, mortality, and extended hospital stay. Furthermore, increased healthcare costs are attributable to SSI.<sup>[3]</sup>

Several perioperative preventive measures have been implemented to minimize the risk of SSI, such as hand washing of the surgical team, antibiotic prophylaxis, skin preparation, and sterile drapes and gowns. Despite these measures healthcare associated infections, especially SSI, remain a challenging problem to surgeons and patients worldwide.<sup>[4]</sup>

Prophylactic (or closed incision) negative pressure wound therapy (pNPWT) denotes the prophylactic use of negative pressure wound therapy (NPWT) to prevent wound complications, specifically SSI. Although NPWT has been used since late 1990s for several purposes, such as open bone fractures,<sup>[5,6]</sup> diabetic ulcers,<sup>[7]</sup> and management of the open abdomen,<sup>[8]</sup> its prophylactic use for primarily closed incisions has only been described a decade ago.<sup>[9]</sup>

Prophylactic NPWT consist of a hermetically sealed system connected to a vacuum pump, which maintains negative pressure on the wound surface. Although several studies on the working mechanism of NPWT have been performed and reviewed,<sup>[10]</sup> there is a lack of preclinical research regarding pNPWT. It has been suggested that by using negative pressure dead space is reduced, tissue proliferation is stimulated, and fluids are removed. Moreover, pNPWT could protect against microorganisms entering from the outside world. Prophylactic NPWT has been suggested as a promising application to reduce SSIs and other wound complications. A few previous reviews have been published<sup>[11,12]</sup> using different methodology, for example, combining randomized controlled trials (RCTs) and observational studies into one analysis. Not many studies on clean-contaminated surgery were included, and none of the previous systematic reviews qualified evidence using Grading of Recommendations Assessment, Development, and Evaluation (GRADE). Our aim was to systematically review the available literature on pNPWT in terms of reducing SSI in all types of surgery. This review was conducted as part of the development of the Global Guidelines for prevention of SSIs commissioned by World Health Organization in Geneva.

## 2. Methods

The PRISMA<sup>[13]</sup> (preferred Reporting Items for Systematic Reviews and Meta-Analyses) and MOOSE<sup>[14]</sup> (meta-analysis of observational studies in epidemiology) guidelines were followed.

### 2.1. Search strategy and selection criteria

References for this review were identified through searches of PubMed, EMBASE (Ovid), the Cochrane Central Register of Controlled Trials (CENTRAL), CINAHL, and World Health Organization for articles published from January 1990 to October 7, 2015 by use of the terms “surgical site infection,” “negative pressure wound therapy,” and “surgical procedure.” The complete search is included in appendix A, <http://links.lww.com/MD/B234>. Two authors (FdV and EW) independently screened all titles and abstracts. RCTs or observational studies

comparing pNPWT with conventional wound dressings in adult patients were included. We chose to include observational studies, because previous systematic reviews did not identify any RCTs in potentially contaminated surgery. Studies had to, at least, report on SSIs or wound infections, either as primary or secondary outcome. Studies using NPWT on an open wound or split skin graft were excluded. We had no language restrictions. References of the included studies were screened for other relevant studies. We only considered full text published studies.

### 2.2. Data extraction

Study characteristics including year of publication, number of patients, types of surgical procedures, duration of prophylactic NPWT, amount of negative pressure, details on standard dressings, and outcome were retrieved from the text.

### 2.3. Quality assessment

Quality of the included studies was assessed with the Cochrane Collaboration’s tool for assessing risk of bias<sup>[15]</sup> for RCTs and with the New-Castle Ottawa scale<sup>[16]</sup> for observational studies. The GRADE methodology was used to assess the quality of the body of retrieved evidence (GRADEpro, Version 20. McMaster University, 2014).

### 2.4. Data synthesis and analysis

Meta-analyses were performed with a random effect model in Review Manager (RevMan), Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014. Heterogeneity of the included studies was evaluated by calculating the  $I^2$  statistic. Subgroup meta-analyses stratified by wound class and type of surgery were performed subsequently.

### 2.5. Ethical approval

Ethical approval was not necessary due to the study design.

## 3. Results

Of the 1238 articles initially identified by the search, we selected 49 for a full text review (Fig. 1). Thirty articles were excluded; 14 studies were not relevant, 14 did not describe pNPWT, 1<sup>[17]</sup> did not report on SSIs as outcome, and 2 studies reported on an overlapping cohort<sup>[18,19]</sup> and were, therefore, combined for the purpose of analysis. Finally, 19 articles describing 21 studies were included. One article described 2 separate randomized controlled trials within one report,<sup>[9]</sup> which were assessed and analyzed separately in present meta-analysis. In another observational study, 2 patient populations (breast and colorectal surgery) were assessed and analyzed separately.<sup>[20]</sup> These were 6 RCTs reported in 5 articles (including 562 patients) and 15 observational studies reported in 14 articles (including 4560 patients).

### 3.1. Study characteristics

We found 6 RCTs,<sup>[6,9,21–23]</sup> 3 prospective observational studies,<sup>[19,20,24]</sup> 10 retrospective observational studies,<sup>[25–34]</sup> and 1 article<sup>[35]</sup> with both retro- and prospective data. In the observational studies, the use of pNPWT was based on the surgeons decision in 6 studies<sup>[19,25,27,30,32,33]</sup> and based on time




Figure 1. Flow-chart of the systematic review.

(before-after) in 8 studies.<sup>[24,25,28,29,31,34–36]</sup> The evidence table with more detailed information is in Table 1.

We found 9 studies on abdominal surgery<sup>[19,26–28,36]</sup> of which 4 involved ventral hernia repair procedures,<sup>[29,30,32,34]</sup> 6 studies in orthopaedic or trauma surgery<sup>[9,21,22,25,33]</sup> 2 studies in cardiac surgery,<sup>[24,35]</sup> and 2 studies in vascular surgery. One study<sup>[20]</sup> included both abdominal and breast surgery. Apart from 1 study, all RCTs were performed in clean surgery. The other study involved patients of which more than 90% underwent lower extremity amputation due to chronic wounds. The negative pressure devices were set between 75 and 125 mm Hg, and the length of negative pressure varied between 24 hours till 7 days postoperative. Either dry gauze, conventional-, occlusive-, or absorbent dressings were used in the control group.

The definition for SSI differed between studies. Nine studies used the definition described by the Centers for Disease Control and Prevention,<sup>[37]</sup> 4 used clinical signs and symptoms as a

criteria for SSI, and 2 other, both retrospective studies, scored for SSI in case any treatment was necessary. Follow-up time varied considerably but was, if reported, always at least 30 days. The risk of bias scoring is provided in appendix A, <http://links.lww.com/MD/B234>.

### 3.2. Prophylactic negative pressure wound therapy versus conventional wound dressings

The meta-analysis of 6 RCTs<sup>[6,9,21–23]</sup> showed a significant benefit of pNPWT over conventional dressings in regard to reducing the risk of SSI (odds ratio [OR], 0.56; 95% confidence interval [CI], 0.32–0.96,  $I^2$  0%). The meta-analysis of 15 observational studies<sup>[19,20,24–35]</sup> showed a significant benefit of pNPWT over conventional dressings in regard to reducing the risk of SSI (OR, 0.30; 95% CI, 0.22–0.42,  $I^2$  18%) (Figs. 2 and 3). The funnel plots of both meta-analyses are included in

**Table 1**  
**Evidence table.**

Study + year	Design	Type of surgery	Population	SSI definition	pNPWT	Conventional dressing	Antibiotic prophylaxis	Outcome (SSI/wound-infection)
<i>RCT</i>								
Gillespie BM et al (2015) <sup>[21]</sup>	RCT	Primary hip arthroplasty	70	CDC	PICO	Hydrocolloid	NR	<i>t</i> : 2/35, <i>C</i> : 3/35, <i>P</i> : 0.65, <i>RR</i> : 0.67, 95% <i>CI</i> : 0.1–3.7
Howell RD (2011) <sup>[22]</sup>	RCT	Total knee arthroplasty	51/60 <sup>§</sup>	NR	VAC	Dry gauze dressing	Yes	<i>t</i> : 1/24, <i>C</i> : 1/36, <i>P</i> <i>OR</i> / <i>CI</i> : NR
Masden D et al (2012) <sup>[23]</sup>	RCT	>90% lower extremity amputation	81	Unclear, but any infection needing an intervention	VAC	Dry sterile dressing	NR	<i>t</i> : 3/44, <i>C</i> : 5/37, <i>P</i> : 0.46
Stannard JP et al (2012) <sup>[6]</sup>	RCT	Lower extremity fracture surgery	263	Combination of clinical signs and symptoms, cultures	VAC	NR dressing	NR	<i>t</i> : 14/141, <i>C</i> : 23/122, <i>P</i> : 0.049
Stannard JP et al (2006) <sup>[9]</sup>	RCT	High energy trauma/hematoma	44	Signs and symptoms consistent with infection	VAC	Pressure dressing	NR	<i>t</i> : 1/13, <i>C</i> : 5/31, <i>P</i> > 0.05
		High energy trauma/healing	44		VAC	NR dressing		<i>t</i> : 3/20, <i>C</i> : 3/24, <i>P</i> > 0.05
<i>Cohort</i>								
Adogwa O et al (2014) <sup>[25]</sup>	RS	Spine surgery	160	CDC	NR/PICO	Sterile dressing	Yes	<i>t</i> : 5/46, <i>C</i> : 17/114, <i>P</i> : 0.04
Blackham AU et al (2013) <sup>[26]</sup>	RS	Colorectal, pancreatic or peritoneal surface surgery	189/191 <sup>§</sup>	CDC	VAC	Standard sterile dressing	Yes	<i>t</i> : 17/104, <i>C</i> : 23/87, <i>P</i> : 0.113, <i>P</i> : 0.065*, <i>OR</i> (95% <i>CI</i> ): 0.48* (0.22–1.05)
Bonds AM et al (2013) <sup>[27]</sup>	RS	Open colorectal surgery	254	CDC	VAC	Occlusive dressing	Yes	<i>t</i> : 4/32, <i>C</i> : 65/222, <i>P</i> : 0.036, <i>OR</i> : 0.290
Chadi SA et al (2014) <sup>[28]</sup>	RS	Abdominoperineal resection	59	CDC	VAC	Gauze	Yes	<i>t</i> : 4/27, <i>C</i> : 13/32, <i>P</i> : 0.01 <i>OR</i> (95% <i>CI</i> ): 0.11 (0.04–0.66) <sup>†</sup>
Condé-Green A et al (2013) <sup>[29]</sup>	RS	Ventral hernia repairs	56	NR	VAC	Dry gauze dressings	NR	<i>t</i> : 1/23, <i>C</i> : 2/33, <i>P</i> <i>OR</i> / <i>CI</i> : NR
Gassman A et al (2015) <sup>[30]</sup>	RS	Abdominal wall reconstruction	61	Clinical findings of incisional cellulitis and leukocytosis	VAC	Dry occlusive dressings	Yes	<i>t</i> : 5/29, <i>C</i> : 17/32, <i>P</i> : 0.01
Grauhan O et al (2013) <sup>[24]</sup>	PS	Cardiac surgery	150	CDC	Prevena	Conventional wound dressings	Yes	<i>t</i> : 3/75, <i>C</i> : 12/75, <i>P</i> : 0.266, <i>OR</i> (95% <i>CI</i> ): 4.57 (1.23–16.94)
Grauhan O et al (2014) <sup>[35]</sup>	PS & RS	Cardiac surgery	3745	NR	Prevena	Conventional wound dressings	NR	<i>t</i> : 3/237, <i>C</i> : 119/3508, <i>P</i> < 0.05, <i>OR</i> : 2.74
Matatov T et al (2013) <sup>[31]</sup>	RS	Vascular surgery (longitudinal or transverse femoral cutdown) with groin incision	90/115 <sup>§</sup>	sign and symptoms + Szilagyi grading	Prevena	Adhesive or absorbent dressing	Yes	<i>t</i> : 3/52, <i>C</i> : 19/63, <i>P</i> : 0.0011
Pauli EM et al (2013) <sup>[32]</sup>	RS	Ventral hernia repairs	119	Unclear, but any infection needing an intervention	VAC	Dry gauze adhesive dressing	Yes	<i>t</i> : 10/49, <i>C</i> : 18/70, <i>P</i> : 0.50
Pellino G et al (2014) <sup>[20]</sup>	PS	Breast surgery	50	CDC	PICO	Absorbent dressing	NR	<i>t</i> : 2/25, <i>C</i> : 9/25, <i>P</i> : 0.04
		Colorectal surgery	50	CDC	PICO	Absorbent dressing	NR	<i>t</i> : 2/25, <i>C</i> : 11/25, <i>P</i> : 0.008

Study + year	Design	Type of surgery	Population	SSI definition	pNPWT	Conventional dressing	Antibiotic prophylaxis	Outcome (SSI/wound-infection)
Reddix RN et al (2010) <sup>[33]</sup>	RS	Acetabular fracture surgery	301	Any infection needing an intervention	VAC	NR	Yes	<i>t</i> : 3/235, <i>C</i> : 4/66, <i>P</i> : 0.0414
Selvaggi et al (2014) <sup>[19]</sup>	PS	Abdominal surgery for Crohn's disease	50	CDC	PICO	Absorbent dressing	Yes	<i>t</i> : 2/25, <i>C</i> : 12/25, <i>P</i> : 0.04
Soares KC et al (2015) <sup>[34]</sup>	RS	Ventral hernia repairs	199	CDC	HVAC	Dry gauze	Yes	<i>t</i> : 10/115, <i>C</i> : 27/84, <i>P</i> : <0.001, <i>P</i> : <0.001 <sup>‡</sup> , OR (95% CI): 0.10 (0.03–0.3) <sup>§</sup>

C=control group, CDC=Centers for Disease Control and Prevention, HVAC=hybrid-VAC, I=intervention group, OR=odds ratio, *P*=*P* value, PICO=single use NPWT device (Smith and Nephew), Prevena incision management system (KCI), PS=prospective, pNPWT=prophylactic negative pressure wound therapy, RCT=random controlled trial, RR=relative risk, RS=retrospective, SSI=surgical site infections, VAC=vacuum assisted closure device (KCI).

\* Adjusted for significant differences between groups using multivariable mixed-effects logistic regression analysis including neo- adjuvant chemotherapy, operation type, wound class, laparoscopic cases, estimated blood loss, patients transfused, number of transfusions, and length of surgery.

<sup>‡</sup> Adjusted OR.

<sup>§</sup> Adjusted using multivariable analysis accounting for age, BMI, ASA class, hernia grade, anesthesia type, operative time, LOS in hospital, wound classification, and mesh location.

<sup>§</sup> Patients/operations.

Appendix B, <http://links.lww.com/MD/B234>. They did not reveal clear indications of publication bias.

### 3.3. Subgroup comparisons

When we stratified observational studies by wound class the meta-analysis of 8 clean-contaminated or contaminated observational studies<sup>[19,20,26–28,30,32,34]</sup> showed a significant benefit of pNPWT over conventional dressings (OR, 0.29; 95% CI, 0.17–0.50) as did the meta-analysis of 8 clean observational studies<sup>[19,20,24,25,29,31,33–35]</sup> (OR, 0.27; 95% CI, 0.17–0.42) (Fig. 4). Two studies<sup>[20,34]</sup> were used in both aforementioned subgroup comparisons as 1 study reported separate results on both clean and clean-contaminated patients and the other study was included as 2 separate cohort studies in our analysis. When we stratified by type of surgery meta-analysis of 4 RCTs in orthopaedic/trauma surgery did not show significant benefit nor harm in regards to reducing the risk of SSI (OR, 0.58; 95% CI, 0.32–1.07). There was only a single RCT in vascular surgery which did not show a significant reduction in SSI, whereas an observational study in vascular surgery did (OR, 0.47; 95% CI, 0.10–2.11 and OR, 0.14; 95% CI, 0.04–0.51, respectively). The meta-analysis of 9 observational studies in abdominal surgery including, besides the 1 study in vascular surgery, ventral hernia repair showed a significant benefit of pNPWT over conventional dressings in regards to reducing SSI (OR, 0.31; 95% CI, 0.19–0.49) as did 2 studies in cardiac surgery (OR, 0.29; 95% CI, 0.12–0.69), and 1 observational study in breast surgery (OR, 0.15; 95% CI: 0.03–0.81). Two observational studies in orthopaedic/trauma surgery showed no significant benefit for pNPWT over conventional dressings (OR, 0.42; 95% CI, 0.13–1.40). These additional subgroup meta-analyses are included in Appendix C, <http://links.lww.com/MD/B234>.

### 3.4. GRADE

Overall evidence was qualified using GRADE for both RCTs and observational studies. Overall, low quality of evidence shows that pNPWT may have benefit when compared to conventional postoperative wound dressings in reducing the risk of SSI. The level of evidence for RCTs was downgraded due to the lack of blinding in outcome assessment in most of the studies and because the optimal information size was not met. The GRADE tables are in Tables 2 and 3.

## 4. Discussion

In this systematic review, a significant benefit was found for pNPWT over conventional wound dressings in terms of reducing SSIs in both RCTs and observational studies (OR, 0.56; 95% CI, 0.32–0.96 and OR, 0.30; 95% CI, 0.22–0.42, respectively). This translates into lowering the SSI rate from 140 to 83 (49–135) per 1000 patients and from 106 to 34 (25–47) per 1000 patients, respectively. These results were consistent in both clean and clean-contaminated procedures and in different types of surgery. However, in stratified analyses, we found no significant benefit for orthopaedic/trauma surgery in both observational and randomized studies, and conflicting data for vascular surgery. Overall quality of evidence, as qualified by GRADE, was low. This means that further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.<sup>[38]</sup>






Figure 2. Overall effect of pNPWT on SSI compared to conventional wound dressings in RCTs. pNPWT = prophylactic negative pressure wound therapy, RCT = randomized controlled trial, SSI = surgical site infections.




Figure 3. Overall effect of pNPWT on SSI compared to conventional wound dressings in observational studies. pNPWT = prophylactic negative pressure wound therapy, SSI = surgical site infections.




Figure 4. Stratification by wound class (all observational studies).

**Table 2****GRADE table RCTs.****Prophylactic negative pressure wound therapy compared to conventional dressings for the prevention of SSI in RCTs**

Quality assessment							Summary of findings		
No. of participants (studies) follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall quality of evidence	Study event rates (%)	Relative effect (95% CI)	Anticipated absolute effects
SSI							With conventional dressings	With prophylactic negative pressure wound therapy	Risk with conventional dressings
562 (6 RCTs)	Serious*	Not serious	Not serious	Serious†	None	⊕⊕⊕⊕ LOW	40/285 (14.0%)	24/277 (8.7%)	OR (95% CI): 0.56 (0.32–0.96)
									Study population
									140 per 1000
									57 fewer per 1000 (91 fewer to 5 fewer)

\* Lack of blinding outcome assessors.

† OIS is not met.

CI = confidence interval, RCT = random controlled trial, SSI = surgical site infections.

**Table 3****GRADE table observational studies.****Prophylactic negative pressure wound therapy compared to conventional dressings for the prevention of SSI in observational studies**

Quality assessment							Summary of findings		
No. of participants (studies) follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall quality of evidence	Study event rates (%)	Relative effect (95% CI)	Anticipated absolute effects
SSI							With conventional dressings	With prophylactic negative pressure wound therapy	Risk with conventional dressings
4560 (15 observational studies)	not serious	not serious	not serious	not serious	none	⊕⊕⊕⊕ LOW	368/3461 (10.6%)	74/1099 (6.7%)	OR (95% CI): 0.30 (0.22–0.42)
									Study population
									106 per 1000
									72 fewer per 1000 (81 fewer to 59 fewer)

CI = confidence interval, SSI = surgical site infections.

Although pNPWT was introduced only 10 years ago, the use of pNPWT has rapidly spread and this application is nowadays used for a variety of wounds. The number of RCTs identified in our literature search is limited and all but 1 RCT were performed following clean, primarily orthopaedic or trauma procedures. We decided to include controlled observational studies because the novelty of the technique is only now generating interest in more accurate RCTs. Moreover, by including observational studies, we included data on potentially contaminated surgery next to clean surgery. In our meta-analyses, we found pNPWT to be more effective in observational studies (OR, 0.30) than RCTs (OR, 0.56). This is expected due to the bias in observational studies where surgeon perceptions, often quite accurate, determine allocation.<sup>[39]</sup> We note that several observational studies included in our review reported a high incidence of SSI in the control group. Most of these studies were performed in abdominal surgery or (complex) ventral hernia repair, surgical procedures prone to SSI.<sup>[40,41]</sup> The inherent selection of high risk wounds prevents extrapolating these results to all types of surgery.

This is the first systematic review and meta-analysis performing separate analyses for observational studies and RCTs and qualifying evidence by the use of GRADE. One previous meta-analysis was performed by Semsarzadeh et al,<sup>[12]</sup> but they combined observational studies and RCTs in their analyses. Additionally, we included 6 recently published studies. Another more recent meta-analysis<sup>[11]</sup> only included RCTs, but they included unpublished data and studies not reporting on the incidence of SSI but only on seroma.

Some limitations of the present study need to be addressed. In the first place, we used (retrospective) observational studies as well as RCTs. Observational studies are well known to involve selection bias and publication bias, as positive results may be more likely to be published than negative results.<sup>[39]</sup> Another limitation was the inclusion of only a small number of patients in most studies, and therefore large confidence intervals within each study. This makes it difficult to estimate the real effect. Although statistical heterogeneity was low ( $I^2$  0% for RCTs and  $I^2$  18% for observational studies), there was variation in protocols used in the included studies in important variables, such as the amount of negative pressure, the duration of pNPWT, and the control dressing. There was also variation in the definition of a SSI used among the included studies and a lack of baseline characteristics influencing the risk of SSI (eg, diabetes or immunosuppression). Therefore, clinical heterogeneity might be higher than statistical heterogeneity reveals.

Although the potential of pNPWT to reduce the incidence of SSI has been shown, limited studies on cost effectiveness are available. The price of pNPWT varies between \$15/d<sup>[42]</sup> and \$495/wk<sup>[43]</sup> depending on a self-made or commercial application. Three studies on cost effectiveness in a gynecologic population have been performed.<sup>[44–46]</sup> Their results show a potential for pNPWT to be cost effective, especially in patient groups with high risk of SSI, such as clean-contaminated operations, or operations performed upon patients with intrinsic risk factors, such as obesity, diabetes, or immunosuppressive medication.

This brings us to the question how useful GRADE is when dealing with new technologies, and even more specific, a new application for a relatively new technology. It remains to be seen whether a significant benefit found in meta-analysis but qualified as low (due to high risk of bias and patient selection) is enough to promote its widespread use. On the other hand, surgeons dealing

with difficult wounds might want to use every option they have to reduce wound infections.

Conclusive data on cost effectiveness are lacking, but pNPWT is most likely to be cost effective in procedures at high risk for SSI due to patients' or operative characteristics. Before widespread use of this application in low-risk as well as high-risk wounds, randomized controlled trials are required to identify the group of patients in whom pNPWT is cost effective. Moreover, the influence of pNPWT on other important wound complications, such as wound dehiscence or seroma, has been shown but needs to be studied more specifically in prospective studies.

## References

- [1] de Lissovoy G, Fraeman K, Hutchins V, et al. Surgical site infection: incidence and impact on hospital utilization and treatment costs. *Am J Infect Control* 2009;37:387–97.
- [2] Magill SS, Edwards JR, Bamberg W, et al. Multistate point-prevalence survey of health care-associated infections. *N Engl J Med* 2014; 370:1198–208.
- [3] Zimlichman E, Henderson D, Tamir O, et al. Health care-associated infections: a meta-analysis of costs and financial impact on the US health care system. *JAMA Intern Med* 2013;173:2039–46.
- [4] World Health Organization (WHO) Report on the Burden of Endemic Health Care-Associated Infection Worldwide. *WHO Libr Cat Data* 2011;40.
- [5] Stannard JP, Volgas DA, Stewart R, et al. Negative pressure wound therapy after severe open fractures: a prospective randomized study. *J Orthop Trauma* 2009;23:552–7.
- [6] Stannard JP, Volgas DA, McGwin G, et al. Incisional negative pressure wound therapy after high-risk lower extremity fractures. *J Orthop Trauma* 2012;26:37–42.
- [7] Rapid T, Service R, Cadth W. Negative pressure wound therapy for managing diabetic foot ulcers: a review of the clinical effectiveness, cost-effectiveness, and guidelines. *CADTH Rapid Response Reports* 2014.
- [8] Roberts DJ, Zygun DA, Grendar J, et al. Negative-pressure wound therapy for critically ill adults with open abdominal wounds. *J Trauma Acute Care Surg* 2012;73:629–39.
- [9] Stannard JP, Robinson JT, Anderson ER, et al. Negative pressure wound therapy to treat hematomas and surgical incisions following high-energy trauma. *J Trauma* 2006;60:1301–6.
- [10] Moues CM, Heule F, Hovius SER. A review of topical negative pressure therapy in wound healing: Sufficient evidence? *Am J Surg* 2011;201: 544–56.
- [11] Hyldig N, Birke-Sorensen H, Kruse M, et al. Meta-analysis of negative-pressure wound therapy for closed surgical incisions. *Br J Surg* 2016; 103:477–86.
- [12] Semsarzadeh NN, Tadisina KK, Maddox J, et al. Closed incision negative-pressure therapy is associated with decreased surgical-site infections. *Plast Reconstr Surg* 2015;136:592–602.
- [13] Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Phys Ther* 2009;89:873–80.
- [14] Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group. *JAMA* 2000; 283:2008–12.
- [15] Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.1.0; 2011.
- [16] Wells G, Shea B, O'Connell D, et al. Newcastle-Ottawa quality assessment scale cohort studies 2014.
- [17] Mark KS, Alger L, Terplan M. Incisional negative pressure therapy to prevent wound complications following cesarean section in morbidly obese women: a pilot study. *Surg Innov* 2014;21:345–9.
- [18] Pellino G, Sciaudone G, Candilio G, et al. Effects of a new pocket device for negative pressure wound therapy on surgical wounds of patients affected with Crohn's disease: a pilot trial. *Surg Innov* 2014;21:204–12.
- [19] Selvaggi F, Pellino G, Sciaudone G, et al. New advances in negative pressure wound therapy (NPWT) for surgical wounds of patients affected with Crohn's disease. *Surg Technol Int* 2014;24:83–9.
- [20] Pellino G, Sciaudone G, Candilio G, et al. Preventive NPWT over closed incisions in general surgery: does age matter? *Int J Surg* 2014;12:S64–8.



- [21] Gillespie BM, Rickard CM, Thalib L, et al. Use of negative-pressure wound dressings to prevent surgical site complications after primary hip arthroplasty: a pilot RCT. *Surg Innov* 2015;22:488–95.
- [22] Howell RD, Hadley S, Strauss E, et al. Blister formation with negative pressure dressings after total knee arthroplasty. *Curr Orthop Pract* 2011;22:176–9.
- [23] Masden D, Goldstein J, Endara M, et al. Negative pressure wound therapy for at-risk surgical closures in patients with multiple comorbidities: a prospective randomized controlled study. *Ann Surg* 2012;255:1043–7.
- [24] Grauhan O, Navasardyan A, Hofmann M, et al. Prevention of poststernotomy wound infections in obese patients by negative pressure wound therapy. *J Thorac Cardiovasc Surg* 2013;145:1387–92.
- [25] Adogwa O, Fatemi P, Perez E, et al. Negative pressure wound therapy reduces incidence of postoperative wound infection and dehiscence after long-segment thoracolumbar spinal fusion: a single institutional experience. *Spine J* 2014;14:2911–7.
- [26] Blackham AU, Farrah JP, McCoy TP, et al. Prevention of surgical site infections in high-risk patients with laparotomy incisions using negative-pressure therapy. *Am J Surg* 2013;205:647–54.
- [27] Bonds AM, Novick TK, Dietert JB, et al. Incisional negative pressure wound therapy significantly reduces surgical site infection in open colorectal surgery. *Dis Colon Rectum* 2013;56:1403–8.
- [28] Chadi SA, Kidane B, Britto K, et al. Incisional negative pressure wound therapy decreases the frequency of postoperative perineal surgical site infections. *Dis Colon Rectum* 2014;57:999–1006.
- [29] Condé-Green A, Chung TL, Holton LH, et al. Incisional negative-pressure wound therapy versus conventional dressings following abdominal wall reconstruction. *Ann Plast Surg* 2012;71:1.
- [30] Gassman A, Mehta A, Bucholdz E, et al. Positive outcomes with negative pressure therapy over primarily closed large abdominal wall reconstruction reduces surgical site infection rates. *Hernia* 2014;19:273–8.
- [31] Matatov T, Reddy KN, Doucet LD, et al. Experience with a new negative pressure incision management system in prevention of groin wound infection in vascular surgery patients. *J Vasc Surg* 2013;3:791–5.
- [32] Pauli EM, Krpata DM, Novitsky YW, et al. Negative pressure therapy for high-risk abdominal wall reconstruction incisions. *Surg Infect (Larchmt)* 2013;14:270–4.
- [33] Reddix RN, Leng XI, Woodall J, et al. The effect of incisional negative pressure therapy on wound complications after acetabular fracture surgery. *J Surg Orthop Adv* 2010;19:91–7.
- [34] Soares KC, Baltodano PA, Hicks CW, et al. Novel wound management system reduction of surgical site morbidity after ventral hernia repairs: a critical analysis. *Am J Surg* 2015;209:324–32.
- [35] Grauhan O, Navasardyan A, Tutkun B, et al. Effect of surgical incision management on wound infections in a poststernotomy patient population. *Int Wound J* 2014;11:6–9.
- [36] Pellino G, Sciaudone G, Candilio G, et al. Effects of a new pocket device for negative pressure wound therapy on surgical wounds of patients affected with Crohn's disease: a pilot trial. *Surg Innov* 2013;21:204–12.
- [37] Mangram AJ, Horan TC, Pearson ML, et al. Guideline for prevention of surgical site infection. *Am J Infect Control* 1999;27:97–134.
- [38] Atkins D, Best D, Briss PA, et al. Grading quality of evidence and strength of recommendations. *Br Med J* 2004;328:1490.
- [39] Kicinski M, Springate DA, Kontopantelis E. Publication bias in meta-analyses from the Cochrane Database of Systematic Reviews. *Stat Med* 2015;34:2781–93.
- [40] Aga E, Keinan-Boker L, Eithan A, et al. Surgical site infections after abdominal surgery: incidence and risk factors. A prospective cohort study. *Infect Dis* 2015;47:761–7.
- [41] Kanters AE, Krpata DM, Blatnik JA, et al. Modified hernia grading scale to stratify surgical site occurrence after open ventral hernia repairs. *J Am Coll Surg* 2012;215:787–93.
- [42] Chaput B, Garrido I, Ebudery H, et al. Low-cost negative-pressure wound therapy using wall vacuum. *Plast Reconstr Surg - Glob Open* 2015;3:e418.
- [43] Prevena™ Incision Management System. Product Monograph 2016. KCI / Acclity.
- [44] Echebiri NC, McDoom MM, Aalto MM, et al. Prophylactic use of negative pressure wound therapy after cesarean delivery. *Obstet Gynecol* 2015;125:299–307.
- [45] Lewis LS, Convery PA, Bolac CS, et al. Cost of care using prophylactic negative pressure wound vacuum on closed laparotomy incisions. *Gynecol Oncol* 2014;132:684–9.
- [46] Tuffaha HW, Gillespie BM, Chaboyer W, et al. Cost-utility analysis of negative pressure wound therapy in high-risk cesarean section wounds. *J Surg Res* 2015;195:612–22.